

- [1] W. M. Haynes, D. R. Lide (Eds.) *CRC Handbook of Chemistry and Physics. A Ready-Reference Book of Chemical and Physical Data*, CRC Press; Taylor & Francis, Boca Raton, Fla., London, **2011**.
- [2] I. G. Vasilyeva, E. S. Vikulova, A. A. Pochtar, N. B. Morozova, *Coatings* **2021**, *11*, 176.
- [3] X. Luo, F. Deng (Eds.) *Micro & Nano Technologies Series*, Elsevier, Amsterdam, Kidlington, Oxford, Cambridge, MA, **2019**.
- [4] M. Fernandes, K. RB Singh, T. Sarkar, P. Singh, R. Pratap Singh, *Adv. Mater. Lett.* **2020**, *11*, 1.
- [5] M. A. Shand, *The Chemistry and Technology of Magnesia*, Wiley-Interscience, Hoboken, N.J, **2006**.
- [6] S. X. Tao, H. W. Chan, H. van der Graaf, *Materials* **2016**, *9*.
- [7] E. G. Jeong, J. H. Kwon, K. S. Kang, S. Y. Jeong, K. C. Choi, *J. Inf. Disp.* **2020**, *21*, 19.
- [8] T. Hatanpää, J. Ihanus, J. Kansikas, I. Mutikainen, M. Ritala, M. Leskelä, *Chem. Mater.* **1999**, *11*, 1846.
- [9] B. W. Wang, J. Choi, H. G. Kim, S. D. Hyun, C. Yoo, S. Kim, H. Lee, C. S. Hwang, *J. Mater. Chem. C* **2021**, *9*, 15359.




Figure 1. Saturation of the precursor pulse using $[\text{Mg}(\text{DMP})_2]$ and H_2O on Si substrates (left) and RBS spectra of the resulting thin films on glassy carbon (right) at 40 °C (green) and 120 °C (blue).