

# Quantum Anomalous Hall Effect in the Magnetic Topological Insulator Thin Films

**Cui-Zu Chang**

*Department of Physics, Pennsylvania State University*

The quantum anomalous Hall (QAH) effect can be considered as the quantum Hall (QH) effect without external magnetic field, which can be realized by time reversal symmetry breaking in a topologically non-trivial system [1, 2]. A QAH system carries spin-polarized dissipationless chiral edge transport channels without the need for external energy input, hence may have huge impact on future electronic and spintronic device applications for ultralow-power consumption. The many decades quest for the experimental realization of QAH phenomenon became a possibility in 2006 with the discovery of topological insulators (TIs). In 2013, the QAH effect was observed in thin films of Cr-doped TI for the first time [3]. Two years later in a near ideal system, V-doped TI, contrary to the negative prediction from first principle calculations, a high-precision QAH quantization with more robust magnetization and a perfectly dissipationless chiral current flow was demonstrated [4]. In this talk, I will introduce the route to the experimental observation of the QAH effect in aforementioned two systems [3, 4], and discuss the demonstration of the dissipationless chiral edge state as well as the origin of the dissipative channels in the QAH state [5]. Finally I will talk about our recent progress on the QAH insulator-Anderson insulator quantum phase transition and its scaling behaviors [6].

[1] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).

[2] R. Yu et al, Science **329**, 61 (2010).

[3] Cui-Zu Chang et al, Science **340**, 167(2013).

[4] Cui-Zu Chang et al, Nature Materials **14**, 473(2015).

[5] Cui-Zu Chang et al, Physics Review Letters **115**, 057206 (2015).

[6] Cui-Zu Chang et al, Physics Review Letters **117**, 126802 (2016).

<sup>+</sup> Author for correspondence: cxc955@psu.edu