

# Monday Evening, December 9, 2024

## Biomaterial Surfaces & Interfaces

### Room Naupaka Salon 5 - Session BI1-MoE

#### Biomaterials/Interfaces - Characterization

Moderator: David G. Castner, University of Washington

5:40pm **BI1-MoE-1 Molecular Structure of Sputtered Species with Large Cluster Ions**, *Jiro Matsuo*, Quantum Science and Engineering Center, Kyoto University, Japan

**INVITED**

Molecular-imaging techniques are now of interest for life science, pharmaceutical and medical applications and organic industrial materials, such as functionalized polymers and organic semiconductors. SIMS (Secondary Ion Mass Spectrometry) is one of the powerful techniques, due to high lateral resolution, high sensitivity and high surface sensitivity. However, ion dose is limited with conventional keV primary ion beam, which destroys molecules of samples with high energy ion impacts. Therefore, SIMS measurement should be completed before organic molecules on the surface are completely damaged. Recently, large cluster ions are utilized for SIMS as a primary ion beam, because there is no limitation in ion dose. It has been demonstrated that large cluster ion beams have a great capability to sputter organic molecules without any residual damage on the surface, because cluster ion beams are equivalently low energy ion beams. Extremely high energy density and multiple collisions are responsible for "cluster effects", which play an important role during their sputtering process of organic molecules. Practical applications, such as organic depth profiling and 3-dimensional molecular analysis in XPS or SIMS have been demonstrated. Nevertheless, high sensitivity is required in all applications, and high sensitivity is becoming an important issue in SIMS technique as well. It is quite important to increase secondary molecular ion yields, which are usually very low (<1E-4). GCIB technique can generate clusters of not only Ar gas but also a variety of gas molecules and various gas molecules are being utilized in the GCIB technique to improve secondary ion yields. Unfortunately, it is still not enough. Most of sputtered species are neutral, but there is no report on molecular structure of sputtered species from organic materials. Due to low ionization yields, SIMS spectra never tell us molecular structure of sputtered species. We have concerned that organic molecules sputtered with large cluster ions are destroyed, or not. Capturing of neutral species and electrospray ionization mass spectrometry (ESI-MS) technique were utilized to explore molecular structure of sputtered neutral species. We have, for the first time, measured the molecular structure of sputtered neutral species and the results clearly show that there are many neutral species maintaining their molecular structure. 10% of neutral particles maintains their molecular structure. If those neutral species are ionized, sensitivity of SIMS would improve by several orders of magnitude.

6:20pm **BI1-MoE-3 GCIB-SIMS Analysis of Skin Cancer Samples**, *John S. Fletcher*, *K. Sjögren Cehajic*, *K. Dimovska Nilsson*, *O. Zaar*, *D. Katsarelis*, *J. Paoli*, *R. Olofsson Bagge*, *N. Neittaanmäki*, University of Gothenburg, Sweden

The use of gas cluster ion beams (GCIBs) for secondary ion mass spectrometry (SIMS) analysis provides softer ejection of biomolecular ions and has created opportunities for meeting the challenges of clinical researchers who require chemical specific imaging of different sample type from cells to tissue biopsies. Here we use a J105 Buncher-ToF SIMS instrument (Ionoptika Ltd, UK) to perform in situ lipidomics of skin cancer samples. GCIB-SIMS analysis enabled detailed spatial-lipidomics that could be directly correlated with conventional histopathological analysis of consecutive H&E slides. Here we present work where melanoma cancer samples were the target in order to investigate the chemical changes associated with disease progression and also to investigate if different metastatic pathways could be distinguished based on the chemical signature of the tumours. Primary tumours were analysed along with "healthy/normal" skin from the same subject along with metastatic tumour samples that had spread via either the lymphatic system or through the blood. Significant differences in the lipid profiles were found in primary compared to metastatic melanomas, notably an increase in phosphatidylethanolamine lipids relative to phosphatidylinositol lipids and an increase in GM3 gangliosides in the metastatic samples. Furthermore, analysis of the data from in-transit versus distant metastases samples highlighted that specific glycerophospholipids, and a difference in the long versus shorter chain GM3 gangliosides, discriminated the metastatic routes. The data is also compared to other skin cancer samples including such as aggressive basal cell carcinoma. Challenges related to data processing and spectral annotation are also discussed.

6:40pm **BI1-MoE-4 Depth Correction of 3D SIMS Depth Profiling Images of Biomaterials Using Only Secondary Ion Signal Intensities**, *M. Brunet*, *B. Gorman*, *Mary Kraft*, University of Illinois Urbana-Champaign

We have developed a depth correction strategy for three-dimensional (3D) SIMS depth profiling images of biomaterials that solely employs secondary ion signal intensity. In this approach, the secondary ion images that were collected during depth profiling are used to create a model of the sample's morphology at the time that each depth profiling image was acquired. Then these models of the sample's morphology are used to shift the voxels in the 3D image to the correct z-position. Comparison of the morphology models created using the secondary ions and the secondary ion images the usage of secondary ion signals with high intensities tends to produce more accurate morphology models. However, even 3D SIMS images that were depth corrected using secondary ions with relatively low intensities were more accurate than uncorrected 3D SIMS depth profiling images. This ability to use secondary ion images to depth correct 3D SIMS depth profiling images in the absence of correlated measurements of sample topography or knowledge of sputter rate expands the range of SIMS depth profiling data sets that may not be depth corrected.

7:00pm **BI1-MoE-5 Label-Free High-Resolution Molecular Imaging of Sex Steroid Hormones in Zebrafish by Water Cluster Secondary Ion Mass Spectrometry (Cluster SIMS)**, *Kate McHardy*, *N. Sano*, Ionoptika Ltd., UK; *E. Lau*, *M. Bailey*, University of Surrey, U.K.

Sex steroid hormones are essential biomolecules for vertebrates and are involved in the maintenance of pregnancy, development of secondary sexual characteristics and diseases such as osteoporosis and breast cancer. Visualising the distribution of steroids contributes to further understanding of disease. However, analysis of steroids is difficult; their low polarity leads to poor ionisation efficiency, meaning they need to be derivatised for conventional analyses. Furthermore, the steroid signals overlap with a MALDI matrix background.

Water Cluster SIMS is a high-sensitivity mass spectrometry technique for imaging complex-mixture materials without derivatisation or the use of matrix. We demonstrate imaging of sex steroid hormones in zebrafish (an ideal vertebrate model organism) with a Water Cluster SIMS instrument.

An adult female zebrafish was prepared for this work. It was embedded while fresh in 0.75% HPMC and 0.25% PVP embedding media to facilitate sectioning. The whole block was flash-frozen in a dry-ice and isopropanol bath. The sample was sectioned to 20 µm at -25 °C and thaw-mounted onto a conductive indium-tin-oxide (ITO) coated glass. The section was dried while frozen in a vacuum desiccator, and then directly analysed without any matrix application for the analysis. The Cluster SIMS analyses were then performed with the J105 SIMS Cluster SIMS (Ionoptika Ltd), using a 70 keV (H<sub>2</sub>O)<sub>n</sub> beam, where n is in the range of 15,000-35,000, and also separately with a 40 keV C<sub>60</sub> beam. High-resolution images were acquired with a pixel size of < 1 micron.

Water Cluster SIMS uses a high-energy beam of ionised clusters of water to sputter and ionise molecules from a surface. It is far less damaging and generates far fewer fragment ions than traditional ToF SIMS, but retains many of the benefits of that technology such as high-spatial-resolution imaging. As a result, detailed images of the distribution of sex steroid hormone molecules in the zebrafish are visible. Preliminary data shows that it is possible to map the chemical distribution of steroids in the ovary area. In addition, we also detected lipid ions related to the embryo or oocyte around the ovary area as unique distributions.

## Author Index

**Bold page numbers indicate presenter**

— **B** —

Bailey, M.: BI1-MoE-5, 1  
Brunet, M.: BI1-MoE-4, 1

— **D** —

Dimovska Nilsson, K.: BI1-MoE-3, 1

— **F** —

Fletcher, J.: BI1-MoE-3, **1**

— **G** —

Gorman, B.: BI1-MoE-4, 1

— **K** —

Katasarelias, D.: BI1-MoE-3, 1  
Kraft, M.: BI1-MoE-4, **1**

— **L** —

Lau, E.: BI1-MoE-5, 1

— **M** —

Matsuo, J.: BI1-MoE-1, **1**  
McHardy, K.: BI1-MoE-5, **1**

— **N** —

Neittaanmäki, N.: BI1-MoE-3, 1

— **O** —

Olofsson Bagge, R.: BI1-MoE-3, 1

— **P** —

Paoli, J.: BI1-MoE-3, 1

— **S** —

Sano, N.: BI1-MoE-5, 1  
Sjögren Cehajic, K.: BI1-MoE-3, 1

— **Z** —

Zaar, O.: BI1-MoE-3, 1